

Published in *Annals of Oncology* 22; 1736-1747, 2011 (on-line 27.6.2011)

## **Strategies for subtypes - dealing with the diversity of breast cancer: highlights of the St.Gallen International Expert Consensus on the Primary Therapy of Early Breast Cancer 2011**

A. Goldhirsch<sup>1\*</sup>, W. C. Wood<sup>2</sup>, A. S. Coates<sup>3</sup>, R. D. Gelber<sup>4</sup>, B. Thürlimann<sup>5</sup>, H.-J. Senn<sup>6</sup> & Panel Members\*\*

\* *Correspondence to:* Prof. A. Goldhirsch, International Breast Cancer Study Group, European Institute of Oncology, Via Ripamonti 435, 20141 Milan, Italy. Tel: +39-02-57489439; Fax: +39-02-94379273; E-mail: aron.goldhirsch@ibcs.org

\*\* See appendix 1 for members of the Panel.

**Pre-publication communication to:** Prof. R. D. Gelber, Dept. of Biostatistics and Computational Biology, CLSB 11007, Dana-Farber Cancer Institute, 450 Brookline Ave., Boston, MA 02215, USA. Tel: +1-617-632-3603; Fax: +1-617-632-2444; E-mail: gelber@jimmy.harvard.edu.

Running head: **Primary breast cancer therapy St.Gallen 2011**

Affiliations of authors:

<sup>1</sup>*International Breast Cancer Study Group, Department of Medicine, European Institute of Oncology, Milan, Italy;* <sup>2</sup>*Department of Surgery, Emory University School of Medicine, N. E. Atlanta, GA, USA;* <sup>3</sup>*International Breast Cancer Study Group and University of Sydney, Sydney, NSW, Australia;* <sup>4</sup>*International Breast Cancer Study Group Statistical Center, Dept. of Biostatistics and Computational Biology, Dana-Farber Cancer Institute and Harvard Medical School, Boston, MA, USA;* <sup>5</sup>*Breast Center, Kantonsspital, St. Gallen, Switzerland;* <sup>6</sup>*Tumor and Breast Center ZeTuP, St. Gallen, Switzerland*

### **Abstract**

The 12<sup>th</sup> St.Gallen International Breast Cancer Conference (2011) expert Panel adopted a new approach to the classification of patients for therapeutic purposes based on the recognition of intrinsic biological subtypes within the breast cancer spectrum. For practical purposes, these subtypes may be approximated using clinico-pathological rather than gene expression array criteria.

In general, systemic therapy recommendations follow the subtype classification. Thus, 'Luminal A' disease generally requires only endocrine therapy, which also forms part of the treatment of the 'Luminal B' subtype. Chemotherapy is considered indicated for most patients with 'Luminal

B', 'HER2 positive', and 'Triple negative (ductal)' disease, with the addition of trastuzumab in 'HER2 positive' disease.

Progress was also noted in defining better tolerated local therapies in selected cases without loss of efficacy, such as accelerated radiation therapy and the omission of axillary dissection under defined circumstances.

Broad treatment recommendations are presented, recognizing that detailed treatment decisions need to consider disease extent, host factors, patient preferences and social and economic constraints.

**Key words:** early breast cancer, St.Gallen Consensus, adjuvant therapies, subtypes

## Introduction

It is no longer tenable to consider breast cancer as a single disease. Subtypes can be defined by genetic array testing (1, 2, 3) or approximations to this classification using immunohistochemistry (4, 5, 6, 7). These subtypes have different epidemiological risk factors (8, 9), different natural histories (10, 11, 12) and different responses to systemic and local therapies (13, 14, 15, 16, 17). These differences imply that clinicians managing breast cancer should consider cases within the various distinct subtypes in order to properly assess the relevant evidence and arrive at appropriate therapeutic advice.

## St.Gallen 2011: news and progress

The 12<sup>th</sup> International Breast Cancer Conference in March 2011 brought together some 4,300 participants from 96 countries and a world-wide faculty representing all relevant disciplines. After presentation of recent research findings, a 51 member expert Panel (see appendix 1) considered a number of questions in order to arrive at treatment recommendations for the immediate future. As in previous St.Gallen conferences (18), the Panel was charged with assessing the evidence, but also advising on the basis of expert opinion on those questions where the evidence was ambiguous or lacking. For the first time, this conference included an explicit approach to management of conflicts of interest (see appendix 2).

Evidence was presented to support a less aggressive approach to axillary surgery in defined circumstances, and the use of more convenient, equally effective approaches to radiation therapy. For systemic therapy, the emphasis of this year's consensus was to reach recommendations within each of the biological subtypes, since these already incorporate many of the risk factors and response predictors previously considered separately. Disease extent, host factors, patient preferences, and economic and social factors inevitably impact the choice and delivery of care. In general, the recommendations are intended to guide therapy considerations outside clinical trials in communities with reasonable levels of available

resources, but noting where possible the availability of alternatives which might be only marginally less effective but less expensive.

This report will first review the new findings presented at the meeting (Table 1), and then proceed to summarize the deliberations of the Panel, bringing these together to form broad therapy recommendations.

**Table 1. Recent research findings presented at the 12<sup>th</sup> International Conference on Primary Therapy of Early Breast Cancer and their implications for patient care**

<b>Field or Treatment</b>	<b>Status of research / Implications for patient care</b>
Surgery – axillary nodes	Several studies have underlined the safety of more conservative approaches to the surgery of the axilla. If sentinel lymph nodes are clear, axillary dissection can be omitted (19). The ACOSOG trial Z0011 for patients with a clinically node negative axilla who underwent lumpectomy and tangential whole breast irradiation showed at a median follow up of 6.3 years that axillary dissection can be omitted without adversely affecting prognosis even in the presence of one or two positive sentinel nodes (20).
Radiation therapy – partial breast irradiation	A randomized trial of targeted intra-operative radiotherapy yielded results closely similar to conventional whole breast irradiation (21). It is noteworthy that in this study 14% of the targeted intra-operative radiotherapy group also received external beam radiotherapy and the median follow up in the study is approximately 2.5 years. A single institution series of 1822 patients treated with breast conserving surgery has documented excellent local control with intra-operative electron beam therapy in selected patients (22).
Radiation therapy – abbreviated (hypofractionated or accelerated) whole breast	Long-term results of the Canadian randomized trial <b>in pT1,2 N0 patients largely treated without adjuvant chemotherapy</b> at a median follow up of 12 years show similar locoregional control, survival, tolerability and cosmesis for a 16 fraction regimen compared with a 25 fraction conventionally fractionated whole breast radiotherapy delivered without external beam boost (23). Similar results have been reported from the UK START trial at a median follow up of 6 years using a 15 fraction regimen (24).
PARP-inhibition	In the presence of tumour defects in homologous recombination DNA repair, inhibition of the PARP enzyme system may result in ‘synthetic lethality’ and increased cell kill (25). This is particularly well seen in carriers of BRCA 1 and 2

	<p>mutations. In such patients, single agent PARP inhibitors, such as olaparib, produce substantial tumour responses. Other cases of Triple Negative disease seldom respond to single agent PARP inhibition (26). In such patients DNA disrupting cytotoxic agents are being investigated in combination with PARP inhibitors.</p>
Anti-HER2 therapies	<p>Double inhibition of HER2 by agents with differing mechanisms of action has been shown to be superior to single agent therapy in neoadjuvant studies (27, 28), a concept being tested in the postoperative adjuvant setting in the ongoing ALTO study. The further study of the mechanism of action of trastuzumab has clarified a role for antibody-dependent cell mediated cytotoxicity (29).</p>
Endocrine therapy in postmenopausal patients	<p>Direct comparison between five years of adjuvant exemestane and anastrozole yielded comparable results, suggesting that exemestane provides an alternative aromatase inhibitor for up-front use (30).</p>
Bisphosphonates	<p>Adjuvant use of zoledronic acid did not improve disease-free survival in a broad population in the AZURE trial (31). Subset analysis of this study showed an apparent benefit in postmenopausal patients and no benefit in premenopausal women. By contrast, the ABCSG 12 trial showed a disease-free survival benefit associated with the use of zoledronic acid among premenopausal patients, all of whom received GnRH analogue (32). These data raise the hypothesis that an antitumour effect of bisphosphonates might depend upon a low estrogen environment. This hypothesis remains to be tested in further clinical trials.</p>
Intrinsic breast cancer subtypes	<p>Definition of intrinsic subtypes has proved efficient in defining prognosis for breast cancer patients (33). Currently, there are no data from phase III trials on their role as predictive tools for chemotherapy benefit. Gene expression arrays are reproducible and quantitative, but cost considerations limit their wide availability. An approximation of gene expression array results is now possible using formalin-fixed, paraffin-embedded material (7).</p>
Molecular mechanisms predicting chemotherapeutic response	<p>Proliferative or immune signatures are associated with good chemotherapy response (34, 35, 36, 37). In neoadjuvant therapy, a stromal signature is associated with a reduced response, while a lymphocytic infiltrate predicts for a higher response rate (38, 39).</p>

Multiple targets for successful treatments	The large and growing number of agents targeting specific mutations suggests the eventual need for individual mutational analysis of each tumour to select a combination of agents to block multiple pathways (40).
Overcoming resistance to endocrine therapies	An improved understanding of the mechanisms of endocrine therapy resistance includes the role of growth factors, integrins, stress kinases, and molecular pathways including PI3K/AKT and MEK/MAPK (41). Overcoming endocrine therapy resistance may, therefore, require inhibition of multiple escape pathways selected by biopsy of resistant tumours to confirm the mechanisms of resistance which are active in each.
Treatment of germline genetic predisposition	Of the 394 genes which have been causally implicated in human cancer, some 10% are transmitted in the germline leading to increased susceptibility (42). Of these the BRCA1 and BRCA2 have been best studied, but others include TP53, PTEN, and CDH1, all of which can increase the risk of breast cancer. BRCA1- and BRCA2-associated breast cancer is sensitive to cross-linking agents such as cisplatin (43), but data from randomized comparisons with standard chemotherapy agents are awaited.
Host factors and cancer risk	Host factors including obesity, and hyperinsulinaemia are associated with increased risk of breast cancer and recurrence of breast cancer. Retrospective studies indicate that diabetic patients receiving metformin have a lower incidence of cancer compared to diabetics not receiving this agent (44).
Vitamins and anti-oxidants	Treatment with beta carotene, vitamin A, and vitamin E may increase mortality. The potential role of vitamin C and selenium on mortality remains inconclusive (45). Fenretinide showed reduced breast cancer incidence in young women (46). The relationship between vitamin D levels and breast cancer risk or prognosis is controversial (46).
Endocrine effects of cytotoxic drugs	A recent analysis of the NSABP B-30 (47) confirmed previous observations from IBCSG 13-93 (48) that amenorrhea following chemotherapy was associated with substantial benefit in disease-free survival. On re-analysis of the NSABP trial using the landmark method, as in the IBCSG study, this effect was limited to the subset of patients with oestrogen receptor positive disease (49).

<p>Modulation of angiogenesis</p>	<p>The early promise of the use of bevacizumab in metastatic breast cancer seen in the E2100 study has not translated into a survival benefit in subsequent studies: a synthesis of these results suggests no overall survival benefit (50). This has led the US Food and Drug Administration to reconsider its accelerated approval of bevacizumab in breast cancer.</p> <p>Studies of lipotransfer have demonstrated the potential for a stromal interaction to stimulate vessel formation, raising the possibility that obesity might have an adverse prognostic impact in cancer patients via a similar stromal interaction (51).</p>
<p>Stem cells</p>	<p>Studies of mammary stem cells suggest a synergistic role for progesterone and RANK ligand in tumour formation (52). This raises the possibility of an additional mechanism of action of clinically available RANK ligand antagonists such as denosumab (53).</p> <p>Further studies of mouse mammary stem cells demonstrated that they are highly responsive to steroid hormone signalling though they lack both estrogen and progesterone receptors. This is thought to be mediated through paracrine signalling involving RANK ligand (54).</p>
<p>Micro RNAs and their influence on tumour growth and inhibition</p>	<p>Micro RNAs are involved in different biopathological features of breast cancer. MER221 and MER222 are involved in resistance and response to endocrine agents, while MER205 is an oncosuppressor able to interfere with response to tyrosine kinase inhibitors of the HER family (55).</p>
<p>Immunity and autoimmunity</p>	<p>Tumour infiltrating regulatory T cells stimulate mammary cancer metastases through receptor activator (RANKL-RANK) signalling (56). Tumour FOXP3<sup>+</sup> T<sub>reg</sub> cells are a major source of RANKL, which stimulates the metastatic progression of HER2-positive, RANK expressing breast cancer cells (52).</p>
<p>Gene based testing</p>	<p>The commercial scores from assays such as Oncotype DX<sup>®</sup> (57) and MammaPrint<sup>®</sup> (58) have been used to determine prognosis. Oncotype DX<sup>®</sup> has been shown to predict chemotherapy benefit among patients with hormone receptor positive disease. An interesting STEPP analysis (59) from the adjuvant trastuzumab NSABP B-31 trial examined the degree of HER2 mRNA expression and corresponding trastuzumab benefit separately for patients with ER positive and ER negative disease. The striking finding was that among patients with ER positive disease, trastuzumab benefit in terms of 8-year disease-free survival was entirely confined to those with the higher levels of HER2 mRNA expression. In contrast, patients with ER negative disease derived some benefit from trastuzumab at all levels of mRNA expression, though the quantitative</p>

	benefit was greater among those with higher levels of HER2 (60).
Timing of adjuvant trastuzumab	The North Central Cancer Treatment Group adjuvant trastuzumab study (N9831) included a randomization between trastuzumab administered either concurrently with or following chemotherapy. Analysis presented at the SABCS 2009 suggested a superior disease-free survival with concurrent administration (61).
Targeted therapy in the neoadjuvant setting	The NOAH study (62) showed clear improvement in breast pathological complete remission (pCR) rate and event-free survival at 3 years with neoadjuvant trastuzumab for patients with HER2-positive disease.
Anti-HER2 therapy without chemotherapy	Studies in metastatic breast cancer and in the neoadjuvant setting have demonstrated activity of trastuzumab and other anti-HER2 agents without chemotherapy (63) albeit usually less than the activity seen for the combination with chemotherapy. There are no corresponding data in the adjuvant setting. However, it may be logical to propose that anti-HER2 therapy, alone or with endocrine therapy if appropriate, may be effective in patients who for various reasons cannot receive cytotoxic therapy (64, 65).
Neoadjuvant platinum in Triple negative (ductal) disease	Triple negative breast cancer includes cases susceptible to DNA damaging agents such as cisplatin. Neoadjuvant studies including cisplatin have produced pCR rates between 22 and 40 percent among unselected triple negative cases (66, 67), while 10 of 12 cases with BRCA1 mutations achieved pCR with single agent cisplatin (68).
Endpoints in neoadjuvant therapy	Failure to achieve pCR among patients with rapidly proliferating tumours identifies a group with a poor prognosis which may be suitable for early trials of investigational agents (11).
Patients with small tumours in the absence of other risk factors	A historical cohort of patients, who did not receive adjuvant systemic therapy in the Danish Breast Cancer Group, were compared with the general Danish population to ascertain mortality ratios associated with the diagnosis of breast cancer. In the absence of other risk factors, patients aged 50 years and older with small (1-10 mm) breast cancers had a risk of death comparable to the background population. By contrast, younger patients with similar tumours had a significantly higher risk of death than the unaffected population (69).

Young patients with endocrine responsive disease	The ABCSG Trial 12 shows that premenopausal women with endocrine responsive disease who receive ovarian function suppression plus either tamoxifen or anastrozole continue to experience a low risk of relapse (32).
Older patients and systemic chemotherapy	The EBCTCG reported similar benefit to systemic chemotherapy in all age groups with ER poor disease (70). The CALGB 49907 study showed inferior results for single agent chemotherapy compared to standard first generation combination regimens (71). The SWOG 8814 trial demonstrated an overall benefit to CAF followed by tamoxifen versus tamoxifen alone in postmenopausal patients with endocrine responsive disease (72), though this was seen primarily among those with adverse biologic features such as quantitatively lower ER levels, involvement of 4 or more lymph nodes, or high 21 gene RS (14).
Special histological types of breast cancer	Review of special histological types in a large institutional series suggested that endocrine responsive types such as tubular and cribriform carcinomas may be suitable for observation without therapy or for endocrine therapy alone. Rare variants of lobular carcinomas (e.g., pleomorphic) and apocrine carcinomas require treatment according to their biological features in a manner analogous to that used for ductal carcinoma. The heterogeneous 'Triple Negative' subtype includes adenoid cystic, juvenile secretory (good prognosis), medullary (intermediate prognosis), and metaplastic (either low-grade, with good prognosis, or high-grade, with poor prognosis) carcinomas, for which no generalizations can be proposed (73).

## Local therapies

New results from clinical trials supported the safety of omitting axillary dissection not only in patients with a negative sentinel node biopsy (19) but also in patients with a clinically node-negative axilla but pathological macrometastatic involvement of one or two sentinel nodes in the context of breast conserving surgery with tangential field radiation therapy (20). This continues a trend of reduced surgical extent without loss of efficacy, which dates back to the breast conserving approaches pioneered by Veronesi (74) and Fisher (75).

Similarly, recent studies in radiation therapy have demonstrated the safety and efficacy of abbreviated schedules for improved patient convenience and the use of partial breast irradiation under certain defined circumstances. These findings are summarized in Table 1.



## Breast cancer subtypes

Analysis of gene expression arrays has resulted in the recognition of several fundamentally different subtypes of breast cancer (1). Because it is not always feasible to obtain gene expression array information, a simplified classification, closely following that proposed by Cheang et al. (7), has been adopted as a useful shorthand. Subtypes defined by clinico-pathological criteria are similar to but not identical to intrinsic subtypes, and represent a convenient approximation. As summarized in Table 2, this approach uses immunohistochemical definition of oestrogen and progesterone receptor, the detection of overexpression and/or amplification of the HER2 oncogene, and Ki-67 labelling index, a marker of cell proliferation, as the means of identifying tumour subtypes.

**Table 2. Surrogate definitions of intrinsic subtypes of breast cancer (4, 7)**

Intrinsic Subtype (1)	Clinico-pathologic definition	Notes
Luminal A	<b>'Luminal A'</b> ER and/or PgR positive(76) HER2 negative (77) Ki-67 low (<14%)*	This cut-point for Ki-67 labelling index was established by comparison with PAM50 intrinsic subtyping (7). Local quality control of Ki-67 staining is important.
Luminal B**	<b>'Luminal B (HER2 negative)'</b> ER and/or PgR positive HER2 negative Ki-67 high  <hr/> <b>'Luminal B (HER2 positive)'</b> ER and/or PgR positive Any Ki-67 HER2 over-expressed or amplified	Genes indicative of higher proliferation are markers of poor prognosis in multiple genetic assays (78). If reliable Ki-67 measurement is not available, some alternative assessment of tumour proliferation such as grade may be used to distinguish between 'Luminal A' and 'Luminal B (HER2 negative)'.  <hr/> Both endocrine and anti-HER2 therapy may be indicated.
Erb-B2 overexpression	<b>'HER2 positive (non luminal)'</b> HER2 over-expressed or amplified ER and PgR absent	
'Basal-like'	<b>'Triple negative (ductal)'</b> ER and PgR absent HER2 negative	Approximately 80% overlap between 'triple negative' and intrinsic 'basal-like' subtype but 'triple negative' also includes some special histological types

		<p>such as (typical) medullary and adenoid cystic carcinoma with low risks of distant recurrence.</p> <p>Staining for basal keratins (79) although shown to aid selection of true basal-like tumours, is considered insufficiently reproducible for general use.</p>
--	--	--

\*This cut-point is derived from comparison with gene array data as a prognostic factor [7]. Optimal cut-points in Ki-67 labelling index for prediction of efficacy of endocrine or cytotoxic therapy may vary.

\*\*Some cases over-express both luminal and HER2 genes.

Clearly, this clinico-pathological classification requires the availability of reliable measurements of its individual components. Guidelines have been published for oestrogen and progesterone receptor determination (76) and for the detection of HER2 positivity (77). For clinical decision making, the Panel supported using the US FDA definition of HER2 positivity based on the eligibility criteria for HER2 status determination from the pivotal clinical trials (80, 81). It was noted that clarifications to the ASCO/CAP guidelines were in preparation, and these have subsequently been published (82). Ki-67 labelling index presents more substantial challenges, but important guidelines for this test are under development (7, 83, 84, 85). In the proposed classification, Ki-67 labelling index is chiefly important in the distinction between ‘Luminal A’ and ‘Luminal B (HER2 negative)’ subtypes. If reliable Ki-67 labelling index assessment is not available some alternative measure of proliferation such as a histological grade may be used in making this distinction.

## Panel deliberations

More than 100 questions were circulated and agreed among Panel members before the meeting. These were presented during the final session of the conference. Panel members had the opportunity to comment, and then voted electronically either yes or no on each question, with the option to abstain if they felt uninformed or conflicted. The detailed votes are not presented here: rather, verbal descriptions of the extent of agreement or disagreement are given in the following sections.

### Axillary surgery

The Panel was clearly of the view that the routine use of immunohistochemistry to look for low volume metastatic disease in sentinel nodes was not indicated, since metastases shown only by immunohistochemistry would not alter management. Furthermore, isolated tumour cells, and

even metastases up to 2 mm (micrometastases) in a single sentinel node, were not considered to constitute an indication for axillary dissection regardless of the type of breast surgery performed. The Panel accepted the option of omitting axillary dissection for macrometastases in the context of lumpectomy and radiation therapy for patients with clinically node-negative disease and 1-2 positive sentinel lymph nodes as reported from ACOSOG trial Z0011 with a median follow-up of 6.3 years (20). The Panel, however, was very clear that this practice, based on a specific clinical trial setting, should not be extended more generally, such as to patients undergoing mastectomy, those who will not receive whole breast tangential field radiation therapy, those with involvement of more than two sentinel nodes, and patients receiving neoadjuvant therapy.

### **Radiation therapy**

The Panel considered accelerated whole breast radiotherapy to be an acceptable option in select patients: in particular, the Panel was divided about the use of this approach in the presence of extensive vascular invasion.

Partial breast irradiation (PBI) as definitive treatment in selected patients was supported by almost half of the Panel and by a strong majority for patients above the age of 70. There was considerable uncertainty about its use in lymphoma survivors who had previously undergone mantle field irradiation, where out-of-quadrant second cancers risks are considerable and for any patient groups different from the current eligible population in PBI trials. The Panel was generally accepting of PBI as an alternative to conventional external beam boost to the tumour bed.

Post-mastectomy radiation therapy was strongly supported for patients with 4 or more involved axillary lymph nodes. While not in general favouring irradiation for those with lesser nodal involvement, the Panel by a slim majority favoured post-mastectomy radiation for patients younger than 45 with 1 to 3 positive nodes and for patients at any age with extensive vascular invasion in 2 or more blocks in conjunction with 1 to 3 positive nodes.

A majority of the Panel supported radiation after complete excision of DCIS, but was prepared to countenance its omission for some elderly patients and those with low grade, low risk DCIS.

### **Definition of biological subtypes**

The Panel strongly supported the clinico-pathological determination of ER, PgR, HER2 and Ki-67 as useful for defining subtypes, but did not support the incorporation of tests for cytokeratin 5/6 or EGFR/HER1 for the determination of 'Basal-like' tumours for clinical decision making. The endorsed clinico-pathological criteria define a convenient alternative to formal subtyping and are likely to be refined in the future. The Panel did not require multi-gene array definition of tumour subtype, although there was acceptance of such assays for certain indications (see below). However, the Panel did recommend that the clinico-pathological markers described above were generally sufficient to guide therapeutic choices.

## Selection of endocrine therapy in premenopausal women

The Panel accepted tamoxifen alone or ovarian function suppression plus tamoxifen as reasonable, though expressing a preference for tamoxifen alone. In patients with a contraindication to tamoxifen, ovarian function suppression alone was accepted as a treatment, while the combination of ovarian function suppression plus an aromatase inhibitor was also considered reasonable.

## Selection of endocrine therapy in postmenopausal women

The Panel was exactly equally divided about whether all postmenopausal patients should receive an aromatase inhibitor (if available and not contraindicated) at some point in treatment, but was more supportive of aromatase inhibitors in the presence of involved lymph nodes. A large majority felt that selected patients could be treated with tamoxifen alone, and that patients could be switched to tamoxifen if intolerant to aromatase inhibitors. The Panel stressed the need to ensure that patients receiving an aromatase inhibitor were indeed postmenopausal, whether by clinical or biochemical criteria.

The Panel considered that 5 years of an aromatase inhibitor was a sufficient duration and a majority opposed extension even in the presence of node-positive disease or among younger postmenopausal patients (less than 55 years of age). The Panel was almost unanimous in rejecting CYP2D6 testing to dictate choice of endocrine therapy type.

## Chemotherapy

The Panel agreed that factors arguing for the inclusion of chemotherapy were high histological grade, high proliferation as measured by Ki-67, low hormone receptor status, positive HER2 status, and 'Triple negative' status in invasive ductal carcinoma of usual forms. These factors are largely captured in the tumour subtype definitions summarized in Table 2. There was a lack of complete consensus on the threshold indication for inclusion of chemotherapy for patients with 'Luminal A' or 'Luminal B (HER2 negative)' disease. In terms of disease extent, the Panel did not believe that node positivity *per se* was an indication for chemotherapy, though a strong majority would use it if more than 3 lymph nodes were involved.

Several tests are available which define prognosis (57, 58, 86). These may indicate a prognosis so good that the doctor and patient decide that chemotherapy is not required. A strong majority of the Panel agreed that the 21-gene signature (Oncotype DX®) (57) may also be used where available to predict chemotherapy responsiveness in an endocrine responsive cohort where uncertainty remains after consideration of other tests, but the majority agreed that the chemo-predictive properties of the 70-gene signature (MammaPrint®) (58) were not yet sufficiently established. Trials are ongoing to clarify this role for both tests. The majority of the Panel did not support lymphovascular invasion as a sufficient indication for chemotherapy, and less than a quarter of the Panel supported uPA/PAI1 (86) as a predictive marker for the use of chemotherapy.

## Chemotherapy in subtypes

The Panel strongly agreed that the 'Luminal A' subtype was less responsive to chemotherapy; that chemotherapy was less useful in such patients; and that no preferred chemotherapy regimen could be defined for treatment of 'Luminal A' disease.

For 'Luminal B' disease, the Panel considered that both anthracyclines and taxanes should be included in the chemotherapy regimen. While the Panel could not define a single preferred chemotherapy regimen for 'HER2 positive' disease, the majority again favoured the inclusion of both anthracyclines and taxanes. For 'Triple negative' disease of the usual ductal type, the Panel again supported the inclusion of anthracyclines and taxanes and an alkylating agent (typically cyclophosphamide), but did not support the routine use of cisplatin or carboplatin. A slim majority agreed that dose-dense chemotherapy (87) should be considered for such patients, and the Panel was strongly opposed to the inclusion of anti-angiogenic therapies at this time, while noting that further trials are ongoing.

## Trastuzumab

The Panel unanimously supported the use of one year of trastuzumab as standard adjuvant treatment for patients with 'HER2 positive' disease, and the majority were willing to extend this to patients with pT1b, but not pT1a pN0 disease. Trastuzumab administered for less than one year (88) was regarded as suboptimal if a year of therapy was feasible, but better than no trastuzumab if limited resources prevented its full duration use. While awaiting data from the ongoing HERA trial, the Panel did not support continuation of adjuvant trastuzumab beyond one year. While preferring that trastuzumab be initiated concurrently with chemotherapy, the Panel also accepted its sequential use. The Panel did not support the use of trastuzumab without chemotherapy if chemotherapy could be given, but was prepared to countenance such treatment in circumstances where chemotherapy could not be delivered.

## Neoadjuvant cytotoxic therapy

A majority of the Panel considered that neoadjuvant cytotoxic therapy was of value beyond its role in facilitating conservative surgery, and noted the improved prognostic information associated with pathological complete response to such therapy, particularly in patients with 'HER2 positive' and 'Triple negative (ductal)' tumours (89) which may allow earlier change from an ineffective regimen.

The Panel considered that the choice of neoadjuvant chemotherapy should be made on the same basis as applied in the selection of postoperative adjuvant treatments. The Panel supported the incorporation of an anti-HER2 drug in the neoadjuvant therapy of patients with 'HER2 positive' disease, but did not support dual HER2 targeting at this point in time. The Panel did not support cytotoxic neoadjuvant therapy for tumours with low proliferation or high endocrine responsiveness.

## Neoadjuvant endocrine therapy

The Panel was almost unanimous in supporting the use of neoadjuvant endocrine therapy as an option for postmenopausal patients with highly endocrine responsive disease. If given, the Panel considered that such treatment should be continued until maximal response or for a minimum of 4-8 months.

## Bisphosphonates

The Panel did not support the use of bisphosphonates for anti-tumour effect in either premenopausal (32) or postmenopausal (90) patients.

## Male breast cancer

Adjuvant tamoxifen was strongly supported, but only a slim majority would consider aromatase inhibitors in patients with contraindications to tamoxifen, such as thrombosis. The Panel did not support extended endocrine treatment beyond 5 years for male breast cancer. The lack of any evidence on these latter two points was acknowledged.

## Summary of systemic treatment recommendations

The approach to treatment within breast cancer subtypes greatly simplifies the definition of therapy indications since the subtypes themselves incorporate many of the risk and predictive factors used in previous consensus recommendations. The broad recommendations are summarized in Table 3, and essentially indicate endocrine therapy alone for patients with clinico-pathologically classified 'Luminal A' disease (except in defined high-risk cases), chemo-endocrine therapy for 'Luminal B', the addition of anti-HER2 therapy in the presence of HER2 positivity and a reliance on chemotherapy for most patients with 'Triple negative' disease (e.g. those with invasive ductal carcinoma).

**Table 3. Systemic treatment recommendations for subtypes**

<b>'Subtype'</b>	<b>Type of therapy</b>	<b>Notes on therapy</b>
<b>'Luminal A'</b>	Endocrine therapy alone	Few require cytotoxics (e.g. high nodal status or other indicator of risk: see text).
<b>'Luminal B (HER2 negative)'</b>	Endocrine ± cytotoxic therapy	Inclusion and type of cytotoxics may depend on level of endocrine receptor expression, perceived risk and patient preference.
<b>'Luminal B (HER2 positive)'</b>	Cytotoxics + anti-HER2 + endocrine therapy	No data are available to support the omission of cytotoxics in this group.

'HER2 positive (non luminal)'	Cytotoxics + anti-HER2	Patients at very low risk (e.g. pT1a and node negative) may be observed without systemic adjuvant treatment.
<p>'Triple negative (ductal)'</p> <p>'Special histological types'*</p> <p>A. Endocrine responsive</p> <hr/> <p>B. Endocrine non responsive</p>	<p>Cytotoxics</p> <hr/> <p>Endocrine therapy</p> <hr/> <p>Cytotoxics</p>	<hr/> <p>Medullary and adenoid cystic carcinomas may not require any adjuvant cytotoxics (if node negative).</p>

\* Special histological types: Endocrine responsive (cribriform, tubular and mucinous); Endocrine non responsive (apocrine, medullary, adenoid cystic and metaplastic).

## Appendix 1 and acknowledgements

Members of the Panel are listed below. All had a significant input to the discussion and manuscript.

- Matti Aapro, Clinique de Genolier, 3 Route du Muids, 1272 Genolier, Switzerland
- Kathy S. Albain, Loyola University Medical Center, Cardinal Bernardin Cancer Center, Room 109, 2160 S First Ave, Maywood, IL 60153, USA
- José Baselga, Massachusetts General Hospital Cancer Center, 55 Fruit Street, YAW 9, Boston, MA 02114, USA
- Jonas Bergh, Radiumhemmet & Karolinska Oncology, Karolinska Institutet and University Hospital, 171 76 Stockholm, Sweden
- Hervé Bonnefoi, Institut Bergonié Cancer Center, Université de Bordeaux, Bordeaux, France
- Harold Burstein, Dept. of Medical Oncology/Solid Tumor Oncology, Dana-Farber Cancer Institute, 450 Brookline Avenue, Boston, MA 02215, USA
- Monica Castiglione-Gertsch, Oncogynecology Unit, University Hospital, Bd. de la Cluse 30, 1211 Geneva, Switzerland
- Alan S. Coates, International Breast Cancer Study Group and University of Sydney, Sydney, NSW 2006, Australia

- Marco Colleoni, Research Unit Medical Senology, European Institute of Oncology, Via Ripamonti 435, 20141 Milan, Italy
- Alberto Costa, ESO – European School of Oncology, Via del Bollo 4, 20123 Milan, Italy and Breast Unit of Southern Switzerland, Lugano, Switzerland
- Giuseppe Curigliano, Div. of Medical Oncology, European Institute of Oncology, Via Ripamonti 435, 20141 Milan, Italy
- Nancy E. Davidson, University of Pittsburgh Cancer Institute, 5150 Centre Avenue, UPMC Cancer Pavilion, 5<sup>th</sup> Floor, Suite 500, Pittsburgh, PA 15232, USA
- Angelo Di Leo, «Sandro Pitigliani» Medical Oncology Unit, Department of Oncology, Hospital of Prato, Piazza dell'Ospedale, 59100 Prato, Italy
- Bent Ejlersen, Dept. of Oncology, Bldg 4262 Rigshospitalet, 9 Blegdamsvej, 2100 Copenhagen, Denmark
- John F. Forbes, Department of Surgical Oncology, University of Newcastle, Calvary Mater Hospital, ANZ BCTG, Edith Street Waratah, NEWCASTLE 2298 NSW Australia.
- Richard D. Gelber, Dept. of Biostatistics and Computational Biology, Dana-Farber Cancer Institute, 450 Brookline Ave., Boston, MA 02215, USA
- Agnes Glaus, ZeTuP, Center for Tumor Detection, Treatment and Prevention, Rorschacher Strasse 150, 9006 St. Gallen, Switzerland
- John H. Glick, University of Pennsylvania, Abramson Cancer Center, Perelman Center, 3400 Civic Center Blvd, Philadelphia, PA 19104, USA
- Michael Gnant, Medical University of Vienna, Dept. of Surgery, Comprehensive Cancer Center Vienna, Währinger Gürtel 18-20, 1090 Wien, Austria
- Aron Goldhirsch, International Breast Cancer Study Group, Swiss Center for Breast Health, Sant'Anna Clinics, 6924 Sorengo-Lugano, Switzerland, and European Institute of Oncology, Via Ripamonti 435, 20141 Milan, Italy (**Chairman**)
- Pamela Goodwin, Dept. of Medicine, Div. of Clinical Epidemiology, Samuel Lunenfeld Research Institute, Mount Sinai Hospital and Princess Margaret Hospital, University of Toronto, 1284-600 University Avenue, Toronto, Ontario, M5G 1X4, Canada
- Paul E. Goss, MGH Cancer Center, 55 Fruit St., Boston, MA 02114, USA
- Jay R. Harris, Dept. of Radiation Oncology, Dana-Farber Cancer Institute, Brigham and Women's Hospital, Rm. 1622, 44 Binney St., Boston, MA 02115, USA
- Dan F. Hayes, Breast Care Center, University of Michigan, Comprehensive Cancer Center, 1500 East Medical Center Drive, Ann Arbor, MI 48109, USA



- James N. Ingle, Mayo Clinic Cancer Center, Women's Cancer Program, 200 First St. S.W., Rochester, MN 55905, USA
- Jacek Jassem, Medical University of Gdansk, Dept. of Oncology & Radiotherapy, Debinki St. 7, 80-211 Gdansk, Poland
- Zefei Jiang, 307 Hospital, Academy of Military Medical Sciences, 8 Fengtai East Avenue, 100071 Beijing, China
- Per Karlsson, Department of Oncology, Institute of Selected Clinical Sciences, Sahlgrenska Academy, Sahlgrenska University Hospital, 413 45 Göteborg, Sweden
- Manfred Kaufmann, Department of Gynecology, Obstetrics, Breast Unit, J.W. Goethe University, Theodor Stern Kai 7, 60596 Frankfurt a. M., Germany
- Gunter von Minckwitz, GBG Forschungs GmbH, Martin Behaim Strasse 12, 63263 Neu-Isenburg, Germany
- Monica Morrow, Memorial Sloan-Kettering Cancer Center, Evelyn Lauder Breast Center, 300 East 66 St NY NY 10065, USA
- Moise Namer, Medical Oncology, Centre Antoine Lacassagne, 33 Av. de Valombrose, 06189 Nice Cedex 2, France
- Larry Norton, Memorial Sloan-Kettering Cancer Center, Rm. H 901, 205 East 64<sup>th</sup> St., Concourse Level, New York, NY 10021-6007, USA
- C. Kent Osborne, Dan L. Duncan Cancer Center, Baylor College of Medicine, One Baylor Plaza, Houston, TX 77030, USA
- Frédérique Penault-Llorca, Service d'Anatomie Pathologie Moléculaire, Dépt. RIO, Centre Jean Perrin, 58 rue Montalembert, BP 392, 63011 Clermont-Ferrand Cedex 1, France
- Charles M. Perou, Lineberger Comprehensive Cancer Center, Departments of Genetics and Pathology, The University of North Carolina, Chapel Hill, North Carolina 27599, USA
- Martine J. Piccart-Gebhart, Internal Medicine, Oncology, Institut Jules Bordet, Rue Héger-Bordet 1, 1000 Brussels, Belgium
- Kurt Possinger, Universitätsklinikum Charité Campus Mitte, Centrum 14, M.S. Onkologie/Hämatologie, Charitéplatz 1, 10117 Berlin, Germany
- Kathleen I. Pritchard, Sunnybrook Odette Cancer Centre, Ontario Clinical Oncology Group (OCOG), 2075 Bayview Avenue, Toronto, ON M4N 1H6, Canada
- Emiel J.T. Rutgers, The Netherlands Cancer Institute, Dept. of Surgery, Plesmanlaan 121, 1066 CX Amsterdam, The Netherlands

- Vladimir Semiglazov, N.N. Petrov Research Institute of Oncology, 68 Leningradskaya St., Pesochny-2, 197758 St. Petersburg, Russia
- Ian Smith, Breast Unit, Royal Marsden Hospital and Institute of Cancer Research, Fulham Road, London, SW3 6JJ, United Kingdom
- Beat Thürlimann, Breast Center, Kantonsspital St. Gallen, 9007 St. Gallen, Switzerland
- Masakazu Toi, Dept. of Breast Surgery, Kyoto University Hospital, 54 Shogoin-Kawahara cho, Sakyo-ku, Kyoto 606-8507, Japan
- Andrew Tutt, Breast Oncology Unit, Guy's Hospital, King's Health Partners AHSC, London, UK
- Michael Untch, Department of Gynecology and Obstetrics, Multidisciplinary Breast Cancer Center, Helios Klinikum Berlin-Buch, Academic Hospital of the University of Göttingen, Schwanebecker Chaussee 50, 13125 Berlin, Germany
- Giuseppe Viale, Dept. of Pathology, European Institute of Oncology and University of Milan, Via Ripamonti 435, 20141 Milan, Italy
- Toru Watanabe, Department of Medicine, Hamamatsu Oncology Center, 3-6-13 Chuo Naka-Ku, 430-0929 Hamamatsu, Japan
- Nicholas Wilcken, Dept. of Medical Oncology, University of Sydney, Westmead Hospital, Westmead 2145, NSW, Australia
- Eric P. Winer, Breast Oncology Center, Dana-Farber Cancer Institute, 450 Brookline Avenue, Boston, MA, 02115, USA
- William C. Wood, Department of Surgery, Winship Cancer Institute, 1365C Clifton Road, Atlanta, GA 30322, USA (**Chairman**)

The authors thank the Participants in the 12<sup>th</sup> International Conference on Primary Therapy of Early Breast Cancer for many useful remarks and for substantial contributions to the process. We acknowledge the substantial contributions of Mrs. Shari Gelber and Mrs. Sabina Briner. We also thank Professor Umberto Veronesi for his guidance and Dr. Franco Nolè for his thoughtful remarks.

## References

- Perou CM, Sorlie T, Eisen MB et al. Molecular portraits of human breast tumours. Nature 2000; 406: 747-752.
- Prat A, Perou CM. Deconstructing the molecular portraits of breast cancer. Mol Oncol 2011; 5: 5-23.

- Parker JS, Mullins M, Cheang MCU et al. Supervised risk predictor of breast cancer based on intrinsic subtypes. *J Clin Oncol* 2009; 27: 1160-1167.
- Nielsen TO, Hsu FD, Jensen K et al. Immunohistochemical and clinical characterization of the basal-like subtype of invasive breast carcinoma. *Clin Cancer Res* 2004; 10: 5367-5374.
- Blows FM, Driver KE, Schmidt MK et al. Subtyping of breast cancer by immunohistochemistry to investigate a relationship between subtype and short and long term survival: a collaborative analysis of data for 10,159 cases from 12 studies. *PLoS Med* 2010; 7: e1000279.
- Hugh J, Hanson J, Cheang MC et al. Breast Cancer Subtypes and Response to Docetaxel in Node-Positive Breast Cancer: Use of an Immunohistochemical Definition in the BCIRG 001 Trial. *J Clin Oncol* 2009; 27: 1168-1176.
- Cheang MCU, Chia SK, Voduc D et al. Ki67 index, HER2 status, and prognosis of patients with luminal B breast cancer. *J Natl Cancer Inst* 2009 ; 101: 736-750.
- Millikan RC, Newman B, Tse CK et al. Epidemiology of basal-like breast cancer. *Breast Cancer Res Treat* 2008; 109: 123-139.
- Phipps AI, Chlebowski RT, Prentice R et al. Body size, physical activity, and risk of triple-negative and estrogen receptor-positive breast cancer. *Cancer Epidemiol Biomarkers Prev* 2011; 20: 454-463.
- Phipps AI, Buist DS, Malone KE et al. Reproductive history and risk of three breast cancer subtypes defined by three biomarkers. *Cancer Causes Control* 2011; 22: 399-405.
- Liedtke C, Mazouni C, Hess KR et al. Response to neoadjuvant therapy and long-term survival in patients with triple-negative breast cancer. *J Clin Oncol* 2008; 26: 1275-1281.
- Dignam JJ, Dukic V, Anderson SJ et al. Hazard of recurrence and adjuvant treatment effects over time in lymph node-negative breast cancer. *Breast Cancer Res Treat* 2009; 116: 595-602.
- Aebi S, Sun Z, Braun D et al. Differential efficacy of three cycles of CMF followed by tamoxifen in patients with ER-positive and ER-negative tumors: Long-term follow up on IBCSG Trial IX. *Ann Oncol* 2011; e-pub 31 January 2011.
- Albain KS, Barlow WE, Shak S et al. Prognostic and predictive value of the 21-gene recurrence score assay in postmenopausal women with node-positive, oestrogen-receptor-positive breast cancer on chemotherapy: a retrospective analysis of a randomised trial. *Lancet Oncol* 2010; 11: 55-65.
- Nguyen PL, Taghian AG, Katz MS et al. Breast cancer subtype approximated by estrogen receptor, progesterone receptor, and HER-2 is associated with local and distant recurrence after breast-conserving therapy. *J Clin Oncol* 2008; 26: 2373-2378.
- Wo JY, Taghian AG, Nguyen PL et al. The association between biological subtype and isolated regional nodal failure after breast-conserving therapy. *Int J Radiat Oncol Biol Phys* 2010; 77: 188-196.
- Tang G, Shak S, Paik S et al. Comparison of the prognostic and predictive utilities of the 21-gene recurrence score assay and Adjuvant! for women with node-negative, ER-positive breast cancer: results from NSABP B-14 and NSABP B-20. *Breast Cancer Res Treat* 2011; e-pub 11 January 2011.
- Goldhirsch A, Ingle JN, Gelber RD et al. Thresholds for therapies: highlights of the St. Gallen International Expert Consensus on the Primary Therapy of Early Breast Cancer 2009. *Ann Oncol* 2009; 20: 1319-1329.
- Krag DN, Anderson SJ, Julian TB et al. Sentinel-lymph-node resection compared with conventional axillary-lymph-node dissection in clinically node-negative patients with breast cancer: overall survival findings from the NSABP B-32 randomised phase 3 trial. *Lancet Oncol* 2010; 11: 927-933.

- Giuliano AE, Hunt KK, Ballman KV et al. Axillary dissection vs no axillary dissection in women with invasive breast cancer and sentinel node metastasis. *JAMA: The Journal of the American Medical Association* 2011; 305: 569-575.
- Vaidya JS, Joseph DJ, Tobias JS et al. Targeted intraoperative radiotherapy versus whole breast radiotherapy for breast cancer (TARGIT-A trial): an international, prospective, randomised, non-inferiority phase 3 trial. *Lancet* 2010; 376: 91-102.
- Veronesi U, Orecchia R, Luini A et al. Intraoperative radiotherapy during breast conserving surgery: a study on 1,822 cases treated with electrons. *Breast Cancer Res Treat* 2010; 124: 141-151.
- Whelan TJ, Pignol JP, Levine MN et al. Long-term results of hypofractionated radiation therapy for breast cancer. *N Engl J Med* 2010; 362: 513-520.
- The START Trialists' Group. The UK Standardisation of Breast Radiotherapy (START) Trial B of radiotherapy hypofractionation for treatment of early breast cancer: a randomised trial. *Lancet* 2008; 371: 1098-1107.
- Tutt A, Robson M, Garber JE et al. Oral poly(ADP-ribose) polymerase inhibitor olaparib in patients with BRCA1 or BRCA2 mutations and advanced breast cancer: a proof-of-concept trial. *Lancet* 2010; 376: 234-244
- Tutt A. PARP inhibitors – walking them through to adjuvant. *The Breast* 2011; 20 (Suppl 1): S1 (Abstr. S01). (manuscript in preparation).
- Baselga J. First results of the NeoALTTO trial (BIG 01-06/EGF 106903): A phase III, randomized, open label, neoadjuvant study of lapatinib, trastuzumab, and their combination plus paclitaxel in women with HER2-positive primary breast cancer. Presented at the San Antonio Breast Cancer Symposium. San Antonio: Texas, 8-12 December 2010; (Abstr. 291).
- Gianni L. Neoadjuvant pertuzumab (P) and trastuzumab (H): Antitumor and safety analysis of a randomized phase II study ('NeoSphere'). Presented at the San Antonio Breast Cancer Symposium. San Antonio: Texas, 8-12 December 2010; (Abstr. 1000).
- Hudis CA. Trastuzumab--mechanism of action and use in clinical practice. *N Engl J Med* 2007; 357: 39-51.
- Goss P, Ingle J, Martino S et al. Outcomes of women who were premenopausal at diagnosis of early stage breast cancer in the NCIC CTG MA17 Trial. Presented at the San Antonio Breast Cancer Symposium. San Antonio: Texas, 8-12 December 2010; (Abstr. S 1-1).
- Coleman RE. Adjuvant treatment with zoledronic acid in stage II/III breast cancer. The AZURE trial (BIG 01/04). Presented at the San Antonio Breast Cancer Symposium. San Antonio: Texas, 8-12 December 2010; (Abstr. 226).
- Gnant M, Mlineritsch B, Schippinger W et al. Endocrine therapy plus zoledronic acid in premenopausal breast cancer. *N Engl J Med* 2009; 360: 679-691.
- Nielsen TO, Parker JS, Leung S et al. A comparison of PAM50 intrinsic subtyping with immunohistochemistry and clinical prognostic factors in tamoxifen-treated estrogen receptor-positive breast cancer. *Clin Cancer Res* 2010; 16: 5222-5232.
- Desmedt C, Haibe-Kains B, Wirapati P et al. Biological processes associated with breast cancer clinical outcome depend on the molecular subtypes. *Clin Cancer Res* 2008; 14: 5158-5165.
- Sotiriou C, Pusztai L. Gene-expression signatures in breast cancer. *N Engl J Med* 2009; 360: 790-800.
- Iwamoto T, Bianchini G, Booser D et al. Gene pathways associated with prognosis and chemotherapy sensitivity in molecular subtypes of breast cancer. *J Natl Cancer Inst* 2011; 103: 264-272.
- Sotiriou C. Molecular mechanisms which predict response to chemotherapy. *The Breast* 2011; 20 (Suppl 1): S3 (Abstr. S05). (manuscript in preparation).

- Andre F, Berrada N, Desmedt C. Implication of tumor microenvironment in the resistance to chemotherapy in breast cancer patients. *Curr Opin Oncol* 2010; 22: 547-551.
- Denkert C, Loibl S, Noske A et al. Tumor-associated lymphocytes as an independent predictor of response to neoadjuvant chemotherapy in breast cancer. *J Clin Oncol* 2010; 28: 105-113.
- Baselga J. Multiple cellular pathways: Evolving targets for successful treatments. *The Breast* 2011; (manuscript in preparation).
- Osborne CK, Schiff R. Mechanisms of endocrine resistance in breast cancer. *Annu Rev Med* 2011; 62: 233-247.
- Daly MB, Axilbund JE, Buys S et al. Genetic/familial high-risk assessment: breast and ovarian. *J Natl Compr Canc Netw* 2010; 8: 562-594.
- Garber J. Germline genetic predisposition: Treatment options. *The Breast* 2011; (manuscript in preparation).
- DeCensi A, Puntoni M, Goodwin P et al. Metformin and cancer risk in diabetic patients: a systematic review and meta-analysis. *Cancer Prev Res (Phila)* 2010; 3: 1451-1461.
- Bjelakovic G, Nikolova D, Gluud LL et al. Mortality in randomized trials of antioxidant supplements for primary and secondary prevention: systematic review and meta-analysis. *JAMA* 2007; 297: 842-857.
- DeCensi A. The science behind vitamins and natural compounds: Getting the most (prevention) out of it. *The Breast* 2011; (manuscript in preparation).
- Swain SM, Jeong JH, Geyer CE, Jr et al. Longer therapy, iatrogenic amenorrhea, and survival in early breast cancer. *N Engl J Med* 2010; 362: 2053-2065.
- International Breast Cancer Study Group, Colleoni M, Gelber S, Goldhirsch A et al. Tamoxifen after adjuvant chemotherapy for premenopausal women with lymph node-positive breast cancer: International Breast Cancer Study Group Trial 13-93. *J Clin Oncol* 2006; 24: 1332-1341.
- Swain SM, Jeong J-H, Wolmark N. Amenorrhea from breast cancer therapy - not a matter of dose. *N Engl J Med* 2010; 363: 2268-2270.
- Burstein HJ. Bevacizumab for advanced breast cancer: All tied up with a RIBBON? *J Clin Oncol* 2011; e-pub 7 March 2011.
- Lohsiriwat V, Curigliano G, Rietjens M et al. Autologous fat transplantation in patients with breast cancer: "silencing" or "fueling" cancer recurrence? *Breast* 2011; e-pub 4 February 2011.
- Schramek D, Leibbrandt A, Sigl V et al. Osteoclast differentiation factor RANKL controls development of progestin-driven mammary cancer. *Nature* 2010; 468: 98-102.
- Gonzalez-Suarez E, Jacob AP, Jones J et al. RANK ligand mediates progestin-induced mammary epithelial proliferation and carcinogenesis. *Nature* 2010; 468: 103-107.
- Asselin-Labat ML, Vaillant F, Sheridan JM et al. Control of mammary stem cell function by steroid hormone signalling. *Nature* 2010; 465: 798-802.
- Iorio MV, Casalini P, Piovan C et al. microRNA-205 regulates HER3 in human breast cancer. *Cancer Res* 2009; 69: 2195-2200.
- Tan W, Zhang W, Strasner A et al. Tumour-infiltrating regulatory T cells stimulate mammary cancer metastasis through RANKL-RANK signalling. *Nature* 2011; 470: 548-553.
- Paik S, Tang G, Shak S et al. Gene expression and benefit of chemotherapy in women with node-negative, estrogen receptor-positive breast cancer. *J Clin Oncol* 2006; 24: 3726-3734.
- Knauer M, Mook S, Rutgers EJ et al. The predictive value of the 70-gene signature for adjuvant chemotherapy in early breast cancer. *Breast Cancer Res Treat* 2010; 120: 655-661.
- Lazar AA, Cole BF, Bonetti M et al. Evaluation of treatment-effect heterogeneity using biomarkers measured on a continuous scale: subpopulation treatment effect pattern plot. *J Clin Oncol* 2010; 28: 4539-4544.
- Paik S. Is gene array testing to be considered routine now? *The Breast* 2011; 20 (Suppl 1): S9 (Abstr. S23). (manuscript in preparation).

- Perez EA, Suman VJ, Davidson NE et al. Results of chemotherapy alone, with sequential or concurrent addition of trastuzumab in the NCCTG N9831 HER2-positive adjuvant breast cancer trial. Presented at the San Antonio Breast Cancer Symposium. San Antonio: Texas, 9-13 December 2009; (Abstr. 80).
- Gianni L, Eiermann W, Semiglazov V et al. Neoadjuvant chemotherapy with trastuzumab followed by adjuvant trastuzumab versus neoadjuvant chemotherapy alone, in patients with HER2-positive locally advanced breast cancer (the NOAH trial): a randomised controlled superiority trial with a parallel HER2-negative cohort. *Lancet* 2010; 375: 377-384.
- Smith I. Is there a case for anti-HER2 treatment without chemotherapy? *The Breast* 2011; 20 (Suppl 1): S16 (Abstr. S41). (manuscript in preparation).
- Kaufman B, Mackey JR, Clemens MR et al. Trastuzumab plus anastrozole versus anastrozole alone for the treatment of postmenopausal women with human epidermal growth factor receptor 2-positive, hormone receptor-positive metastatic breast cancer: Results from the randomized phase III TAnDEM study. *J Clin Oncol* 2009; 27: 5529-5537.
- Schwartzberg LS, Franco SX, Florance A et al. Lapatinib plus letrozole as first-line therapy for HER-2+ hormone receptor-positive metastatic breast cancer. *Oncologist* 2010; 15: 122-129.
- Silver DP, Richardson AL, Eklund AC et al. Efficacy of neoadjuvant cisplatin in triple-negative breast cancer. *J Clin Oncol* 2010; 28: 1145-1153.
- Torrisi R, Cardillo A, Canello G et al. Phase II trial of combination of pegylated liposomal doxorubicin, cisplatin, and infusional 5-fluorouracil (CCF) plus trastuzumab as preoperative treatment for locally advanced and inflammatory breast cancer. *Clin Breast Cancer* 2010; 10: 483-488.
- Byrski T, Gronwald J, Huzarski T et al. Pathologic complete response rates in young women with BRCA1-positive breast cancers after neoadjuvant chemotherapy. *J Clin Oncol* 2010; 28: 375-379.
- Ejlersen B. Characteristics of breast cancer patients unlikely to benefit from adjuvant therapy. *The Breast* 2011; 20 (Suppl 1): S14 (Abstr. S37). (manuscript in preparation).
- Early Breast Cancer Trialists' Collaborative Group. Adjuvant chemotherapy in oestrogen-receptor-poor breast cancer: patient-level meta-analysis of randomised trials. *Lancet* 2008; 371: 29-40.
- Muss HB, Berry DA, Cirincione CT et al. Adjuvant chemotherapy in older women with early-stage breast cancer. *N Engl J Med* 2009; 360: 2055-2065.
- Albain KS, Barlow WE, Ravdin PM et al. Adjuvant chemotherapy and timing of tamoxifen in postmenopausal patients with endocrine-responsive, node-positive breast cancer: a phase 3, open-label, randomised controlled trial. *Lancet* 2009; 374: 2055-2063.
- Colleoni M. Adjuvant therapies for special types of breast cancer. *The Breast* 2011; 20 (Suppl 1): S15 (Abstr. S40). (manuscript in preparation).
- Veronesi U, Saccozzi R, Del VM et al. Comparing radical mastectomy with quadrantectomy, axillary dissection, and radiotherapy in patients with small cancers of the breast. *N Engl J Med* 1981; 305: 6-11.
- Fisher B, Bauer M, Margolese R et al. Five-year results of a randomized clinical trial comparing total mastectomy and segmental mastectomy with or without radiation in the treatment of breast cancer. *N Engl J Med* 1985; 312: 665-673.
- Hammond ME, Hayes DF, Dowsett M et al. American Society of Clinical Oncology/College of American Pathologists guideline recommendations for immunohistochemical testing of estrogen and progesterone receptors in breast cancer. *J Clin Oncol* 2010; 28: 2784-2795.
- Wolff AC, Hammond ME, Schwartz JN et al. American Society of Clinical Oncology/College of American Pathologists guideline recommendations for human epidermal growth factor receptor 2 testing in breast cancer. *J Clin Oncol* 2007; 25: 118-145.

- Wirapati P, Sotiriou C, Kunkel S et al. Meta-analysis of gene expression profiles in breast cancer: toward a unified understanding of breast cancer subtyping and prognosis signatures. *Breast Cancer Res* 2008; 10: R65.
- Cheang MC, Voduc D, Bajdik C et al. Basal-like breast cancer defined by five biomarkers has superior prognostic value than triple-negative phenotype. *Clin Cancer Res* 2008; 14: 1368-1376.
- Piccart-Gebhart MJ, Procter M, Leyland-Jones B et al. Trastuzumab after adjuvant chemotherapy in HER2-positive breast cancer. *N Engl J Med* 2005; 353: 1659–1672.
- Romond EH, Perez EA, Bryant J, et al. Trastuzumab plus adjuvant chemotherapy for operable HER2-positive breast cancer. *N Engl J Med* 2005; 353: 1673–1684.
- Hammond ME, Hayes DF, Wolff AC. Clinical notice for American Society of Clinical Oncology-College of American Pathologists guideline recommendations on ER/PgR and HER2 testing in breast cancer. *J Clin Oncol* 2011. Published Ahead of Print on April 18, 2011 as 10.1200/JCO.2011.35.2245. The latest version is at <http://jco.ascopubs.org/cgi/doi/10.1200/JCO.2011.35.2245>.
- Viale G, Regan MM, Mastropasqua MG et al. Predictive value of tumor Ki-67 expression in two randomized trials of adjuvant chemoendocrine therapy for node-negative breast cancer. *J Natl Cancer Inst* 2008; 100: 207-212.
- Viale G, Giobbie-Hurder A, Regan MM et al. Prognostic and predictive value of centrally reviewed Ki-67 labeling index in postmenopausal women with endocrine-responsive breast cancer: Results from Breast International Group trial 1-98 comparing adjuvant tamoxifen with letrozole. *J Clin Oncol* 2008; 26: 5569-5575.
- Dowsett M, Nielsen TO, A'Hern R et al. Ki67 in breast cancer: Recommendations from the International Ki67 in Breast Cancer Working Group. (pending publication).
- Janicke F, Schmitt M, Pache L et al. Urokinase (uPA) and its inhibitor PAI-1 are strong and independent prognostic factors in node-negative breast cancer. *Breast Cancer Res Treat* 1993; 24: 195-208.
- Citron ML, Berry DA, Cirincione C et al. Randomized Trial of Dose-Dense Versus Conventionally Scheduled and Sequential Versus Concurrent Combination Chemotherapy as Postoperative Adjuvant Treatment of Node-Positive Primary Breast Cancer: First Report of Intergroup Trial C9741/Cancer and Leukemia Group B Trial 9741. *J Clin Oncol* 2003; 21: 1431-1439.
- Joensuu H, Bono P, Kataja V et al. Fluorouracil, Epirubicin, and Cyclophosphamide With Either Docetaxel or Vinorelbine, With or Without Trastuzumab, As Adjuvant Treatments of Breast Cancer: Final Results of the FinHer Trial. *J Clin Oncol* 2009; 27: 5685-5692.
- Kaufmann M, Hortobagyi GN, Goldhirsch A et al. Recommendations from an international expert panel on the use of neoadjuvant (primary) systemic treatment of operable breast cancer: an update. *J Clin Oncol* 2006; 24: 1940-1949.
- Eidtmann H, de Boer R, Bundred N, et al. Efficacy of zoledronic acid in postmenopausal women with early breast cancer receiving adjuvant letrozole: 36-month results of the ZO-FAST Study. *Ann Oncol* 2010; 21: 2188-2194.